

# 2003 Health and Preventive Practices Status Report

## from Missouri County-level Study



# Carroll County\*\*

	County Prevalence (%)*	Northwest Region Prevalence (%)*	State Prevalence (%)*
<b>G</b> eneral			
Health Status (Fair or Poor)	16.2	18.5	16.9
No Health Coverage	9.1	12.1	12.3
<b>B</b> ehavioral Risk Factors			
Physical Inactivity	29.7	26.3	23.0
Current Smoker	27.1	27.3	26.5
Overweight (BMI 25-29.9)	38.7	38.0	35.3
Obese (BMI ≥ 30)	26.2	23.1	23.4
<b>D</b> isease and Conditions			
High Blood Pressure	34.0	28.4	28.5
High Cholesterol (35 years or older)	41.5	40.2	37.3
Asthma	8.1	9.3	10.7
Diabetes	7.1	6.3	7.2
Activity Limitation	21.4	17.5	17.8
<b>C</b> ancer Screening			
<b>Women</b>			
Never had a Mammogram	18.5	17.0	13.4
No Mammogram or Clinical Breast Exam in Last Year:			
40-49 years of age	29.8	44.1	49.4
50-64 years of age	64.1	45.8	35.9
65 and older	81.6 H	62.3 H	47.4
Never had a Pap Test:			
18-69 years of age	0.0 L#	2.8	5.1
70 and older	6.1	12.6	6.4
No Pap Test in Last Year (18-29)	28.7	21.5	25.8
No Pap Test in Last 3 Years (30-69)	10.2	13.9	13.8
<b>Men and Women 50 Years or Older</b>			
Blood Stool Test:			
Never had a Blood Stool Test	59.6	51.0	57.0
No Blood Stool Test in Last Year	83.2	77.8	77.5
Sigmoidoscopy/Colonoscopy			
Never had a Sigmoidoscopy or Colonoscopy	56.2	58.6 H	49.2
No Sigmoidoscopy or Colonoscopy in Past Five Years	64.1	68.6 H	57.0

\*Prevalence rates are age adjusted to the 2000 U.S. standard population

■ indicates that the prevalence rate is not significantly above or below state prevalence

H indicates that the prevalence rate is significantly above the state prevalence

L indicates that the prevalence rate is significantly below the state prevalence

Unless otherwise noted respondents are 18 years of age or older

\*\* The best indicators for Carroll county were obtained by combining data from Carroll and Johnson counties.

# The number of respondents in this category is 0.